arXiv:1411.3488v1 [cond-mat.stat-mech] 13 Nov 2014

The subordinated processes controlled by a family of subordinators and corresponding Fokker-Planck type equations

Long Shi[1,2], Zu-Guo Yu[1,3*], Hai-Lan Huang[1], Zhi Mao[1] and Ai-Guo Xiao[1]

[1] Hunan Key Laboratory for Computation and Simulation in Science and Engineering and Key Laboratory of Intelligent Computing and Information Processing of Ministry of Education, Xiangtan University, Xiangtan, Hunan 411105, China.

[2] School of Science, Central South University of Forest and Technology, Changsha, Hunan 410004, China.

[3]School of Mathematical Sciences, Queensland University of Technology, GPO Box 2434, Brisbane, Q4001, Australia.

Abstract

In this work, we consider subordinated processes controlled by a family of subordinators which consist of a power function of time variable and a negative power function of $\alpha-$stable random variable. The effect of parameters in the subordinators on the subordinated process is discussed. By suitable variable substitutions and Laplace transform technique, the corresponding fractional Fokker-Planck-type equations are derived. We also compute their mean square displacements in a free force field. By choosing suitable ranges of parameters, the resulting subordinated processes may be subdiffusive, normal diffusive or superdiffusive.

Key words: subordinated process; subordinator; Fokker-Planck-type equation; mean square displacement; variable substitution; Laplace transform

1 Introduction

Anomalous diffusion is one of the most universal phenomena in nature [1-3]. In one dimension, it is characterized by a mean square displacement (MSD) of the form

$$\langle (\Delta x)^2 \rangle \sim Dt^{\alpha} \tag{1}$$

with $\alpha \neq 1$, deviating from the linear dependence on time found for normal diffusion [1]. The coefficient D is generalized diffusion constant. It is called subdiffusion for $0 < \alpha < 1$ and superdiffusion for $\alpha > 1$ [4].

*Corresponding to yuzuguo@aliyun.com

One of the simplest models to describe anomalous diffusion is continuous time random walk (CTRW), introduced by Montroll and Weiss [5]. Let $W(x,t)$ be the probability density function (PDF) of finding a particle (with continuous distribution of step length) at point x at time t, one has [5]

$$W(x,t) = \sum_n f(x,n) g_n(t), \tag{2}$$

where $f(x,n)$ is the PDF to find the particle at point x after n steps and $g_n(t)$ is the probability to make exactly n steps up to time t. Here, the random number of steps n plays the role of operational time governing the evolution of the system.

A continuous realization of Eq.(2) is

$$W(x,t) = \int_0^\infty f(x,\tau) g(\tau,t) d\tau, \tag{3}$$

which has been considered by Fogedby in [6] in terms of a coupled system of Itô stochastic differential equations for position x and time t

$$\frac{dx(\tau)}{d\tau} = F(x) + \Gamma(\tau), \qquad \frac{dt(\tau)}{d\tau} = \eta(\tau), \tag{4}$$

where $F(x)$ is an external force, $\Gamma(\tau)$ and $\eta(\tau)$ are random noise sources that are assumed to be independent. In this frame, the position x and the physical time t are parameterized in terms of the continuous variable τ which may be considered as operational time. The random walk in physical time $Y(t)$ is then given by the combined process $Y(t) = x(s(t))$, where $\tau = s(t)$ is the inverse process to $t(\tau)$ defined as

$$s(t) = inf\{\tau > 0 : t(\tau) > t\}, \tag{5}$$

a continuous analogue of the random number of the steps $n(t)$.

In formula (3) $W(x,t)$, $f(x,\tau)$ and $g(\tau,t)$ are the PDFs of the processes $Y(t)$, $x(\tau)$ and $s(t)$, respectively. From the opinion of subordination, which is originated by Bochner [7], the combined process $Y(t)$ can be said to be subordinated to parent process $x(\tau)$ using the operational time $\tau = s(t)$. The process $s(t)$ is also called subordinator. It is worth to note that the continuous time random processes in phase (position-velocity) space based on the subordination were discussed in Refs. [8-10].

In system (4), the most popular choice on random noise sources $\Gamma(\tau), \eta(\tau)$ is taken $\Gamma(\tau)$ as a Gaussian noise and $\eta(\tau)$ as a fully skew $\alpha-$stable Lévy noise (i.e. $s(t)$ is an inverse $\alpha-$stable subordinator). In recent years, the system of coupled Langevin equations (4) with inverse $\alpha-$stable subordinator is becoming a hot topic [11-25]. The inverse tempered $\alpha-$stable subordinator and the infinitely divisible inverse subordinators were also considered [26-34].

However, one cannot obtain directly characteristics of subordinator $s(t)$ from the definition (5). In [35], Meerschaert and Scheffler showed that the $\alpha-$self-similar process $E(t)$ has the same distribution as the process $t^\alpha Z^{-\alpha}$, where Z is an $\alpha-$stable random variable, i.e. $\langle e^{-uZ} \rangle = e^{-u^\alpha}$. Motivated by this, we assume that the subordinator $s(t)$ has the following three forms $s(t) = t^\alpha Z^{-\alpha}$, $s(t) = t^\beta Z^{-\alpha}$ or $s(t) = t^\beta Z^{-\beta}$ and then discuss the resulting subordinated processes $x(s(t))$.

This paper is organized as follows. In section 2, we introduce the subordinated process controlled by subordinator $s(t) = t^{\alpha} Z^{-\alpha} (0 < \alpha < 1)$ and connect the subordinated process $x(s(t))$ with a time-fractional Fokker-Planck equation (FPE). In section 3, we discuss the subordinated process $x(s(t))$ with $s(t) = t^{\beta} Z^{-\alpha} (0 < \alpha < 1, 0 < \beta < 2)$. The subordinated process $x(s(t))$ with $s(t) = t^{\beta} Z^{-\beta} (0 < \beta < 2)$ is considered in section 4. The conclusions are presented in section 5.

2 The subordinated process with subordinator $s(t) = t^{\alpha} Z^{-\alpha}$

Assumed that the *parent process* $x(\tau)$ satisfies the Langevin equation

$$\frac{dx(\tau)}{d\tau} = F(x(\tau)) + \Gamma(\tau), \tag{6}$$

where $\Gamma(\tau)$ is a white Gaussian noise with $\langle \Gamma(\tau) \rangle = 0, \langle \Gamma(\tau)\Gamma(\tau') \rangle = 2\delta(\tau - \tau')$. Thus, the PDF $f(x, \tau)$ obeys the following ordinary FPE [36]:

$$\frac{\partial}{\partial \tau} f(x, \tau) = (-\frac{\partial}{\partial x} F(x) + \frac{\partial^2}{\partial x^2}) f(x, \tau) \triangleq L_{FP} f(x, \tau). \tag{7}$$

Now we assume $s(t) = t^{\alpha} Z^{-\alpha} (0 < \alpha < 1)$, where Z is an $\alpha-$stable random variable with the PDF $\psi_{\alpha}(t)$, i.e. the Laplace transform $\langle e^{-uZ} \rangle = \int_0^{\infty} \psi_{\alpha}(t) e^{-ut} dt = e^{-u^{\alpha}}$.

Since

$$P(s(t) \leq \tau) = P(Z \geq \frac{t}{\tau^{1/\alpha}}) = \int_{\frac{t}{\tau^{1/\alpha}}}^{\infty} \psi_{\alpha}(t') dt', \tag{8}$$

we have the PDF $g(\tau, t)$ of subordinator $s(t)$ as

$$g(\tau, t) = \frac{1}{\alpha} \frac{t}{\tau^{1+1/\alpha}} \psi_{\alpha}(\frac{t}{\tau^{1/\alpha}}). \tag{9}$$

Recalling the following Laplace transform formulas

$$\int_0^{\infty} \varphi(at) e^{-ut} dt = \frac{1}{a} \int_0^{\infty} \varphi(t') e^{-\frac{u}{a} t'} dt' = \frac{1}{a} \widetilde{\varphi}(\frac{u}{a}), \quad a > 0, \tag{10}$$

and

$$\int_0^{\infty} t\varphi(t) e^{-ut} dt = -\frac{d}{du} \int_0^{\infty} \varphi(t) e^{-ut} dt = -\frac{d}{du} \widetilde{\varphi}(u), \tag{11}$$

we get the Laplace transform of $g(\tau, t)$ as

$$\widetilde{g}(\tau, u) = u^{\alpha - 1} \exp\{-u^{\alpha} \tau\}. \tag{12}$$

Substituting $\widetilde{g}(\tau, u)$ into Eq.(3), we get the PDF $W(x, t)$ of the subordinated process $Y(t) = x(s(t))$ in Laplace space:

$$\begin{aligned} \widetilde{W}(x, u) &= \int_0^{\infty} f(x, \tau) \widetilde{g}(\tau, u) d\tau \\ &= \int_0^{\infty} f(x, \tau) u^{\alpha - 1} \exp\{-u^{\alpha} \tau\} d\tau \\ &= u^{\alpha - 1} \widetilde{f}(x, u^{\alpha}). \end{aligned} \tag{13}$$

Since $f(x,\tau)$ satisfies Eq.(7), one get in Laplace space

$$u\widetilde{f}(x,u) - f(x,0) = L_{FP}\widetilde{f}(x,u). \tag{14}$$

Assumed that $W(x,0) = f(x,0) = \delta(x)$, from Eq.(13) and Eq.(14), we have

$$\begin{aligned} u\widetilde{W}(x,u) - W(x,0) &= u^{\alpha}\widetilde{f}(x,u^{\alpha}) - f(x,0) \\ &= L_{FP}\widetilde{f}(x,u^{\alpha}) \\ &= u^{1-\alpha}L_{FP}\widetilde{W}(x,u). \end{aligned} \tag{15}$$

Taking Laplace inverse transform for $u \to t$ in Eq.(15), we obtain the following time-fractional FPE [4,37-39]

$$\frac{\partial W(x,t)}{\partial t} =_0 D_t^{1-\alpha} L_{FP} W(x,t), \quad 0 < \alpha < 1, \tag{16}$$

where ${}_0D_t^{1-\alpha}$ denotes the Riemann-Liouville fractional derivative [40], defined by

$${}_0D_t^{1-\alpha} f(t) = \frac{1}{\Gamma(\alpha)}\frac{\partial}{\partial t}\int_0^t (t-s)^{\alpha-1} f(s)ds. \tag{17}$$

Eq.(16) can be also derived from a generalised master equation [41], or from continuous time random walks (CTRWs) [42].

We are also interested in the MSD of the subordinated process $Y(t)$ in free force field. One can compute $\langle x^2\rangle(t)$ easily in terms of Eq.(16) with natural boundary conditions:

$$\langle x^2\rangle(t) = \int_{-\infty}^{\infty} x^2 W(x,t)dx = \frac{2}{\Gamma(\alpha+1)}t^{\alpha}, \quad 0 < \alpha < 1. \tag{18}$$

From (18), we conclude that the proposed subordinated process is subdiffusive.

3 The subordinated process with subordinator $s(t) = t^{\beta}Z^{-\alpha}$

In this section, we assume that $x(\tau)$ satisfies Langevin equation (6) and $s(t)$ has the form $s(t) = t^{\beta}Z^{-\alpha}(0 < \alpha < 1, 0 < \beta < 2)$.

Since

$$P(s(t) \le \tau) = P(Z \ge \frac{t^{\beta/\alpha}}{\tau^{1/\alpha}}) = \int_{\frac{t^{\beta/\alpha}}{\tau^{1/\alpha}}}^{\infty} \psi_{\alpha}(t')dt', \tag{19}$$

we have

$$g(\tau,t) = \frac{1}{\alpha}\frac{t^{\beta/\alpha}}{\tau^{1+1/\alpha}}\psi_{\alpha}(\frac{t^{\beta/\alpha}}{\tau^{1/\alpha}}). \tag{20}$$

Setting

$$h(\tau,t) = \frac{1}{\alpha}\frac{t}{\tau^{1+1/\alpha}}\psi_{\alpha}(\frac{t}{\tau^{1/\alpha}}), \tag{21}$$

one can easily obtain

$$g(\tau,t) = h(\tau,t_1) \tag{22}$$

with $t_1 = t^{\beta/\alpha}$.

Since

$$W(x,t) = \int_0^\infty f(x,\tau)g(\tau,t)d\tau, \tag{23}$$

we have

$$\begin{aligned} \frac{\partial}{\partial t}W(x,t) &= \int_0^\infty f(x,\tau)\frac{\partial}{\partial t}g(\tau,t)d\tau \\ &= \int_0^\infty f(x,\tau)\frac{\partial h(\tau,t_1)}{\partial t_1}\frac{dt_1}{dt}d\tau \\ &= \frac{\partial}{\partial t_1}\int_0^\infty f(x,\tau)h(\tau,t_1)d\tau\frac{dt_1}{dt}. \end{aligned} \tag{24}$$

Setting $W_1(x,t) = \int_0^\infty f(x,\tau)h(\tau,t)d\tau$, by comparing Eq.(21) with Eq.(9), we obtain that $W_1(x,t)$ satisfies

$$\frac{\partial W_1(x,t)}{\partial t} =_0 D_t^{1-\alpha}L_{FP}W_1(x,t). \tag{25}$$

Substituting Eq.(25) into Eq.(24), we have

$$\begin{aligned} \frac{\partial}{\partial t}W(x,t) &= \frac{\partial}{\partial t_1}W_1(x,t_1)\cdot\frac{dt_1}{dt} \\ &= \frac{1}{\Gamma(\alpha)}\frac{\partial}{\partial t_1}\int_0^{t_1}(t_1-s)^{\alpha-1}L_{FP}W_1(x,s)ds\cdot\frac{dt_1}{dt} \\ &= \frac{1}{\Gamma(\alpha)}\frac{\partial}{\partial t}\int_0^{t_1}(t_1-s)^{\alpha-1}L_{FP}W_1(x,s)ds \\ &= \frac{1}{\Gamma(\alpha)}\frac{\partial}{\partial t}\int_0^{t^{\beta/\alpha}}(t^{\beta/\alpha}-s)^{\alpha-1}L_{FP}W_1(x,s)ds. \end{aligned} \tag{26}$$

After replacing the variable s by $t'^{\beta/\alpha}$ and using the relation $W_1(x,t^{\beta/\alpha}) = W(x,t)$ in Eq.(26), we obtain

$$\begin{aligned} \frac{\partial}{\partial t}W(x,t) &= \frac{\beta}{\Gamma(\alpha+1)}\frac{\partial}{\partial t}\int_0^t t'^{\beta/\alpha-1}(t^{\beta/\alpha}-t'^{\beta/\alpha})^{\alpha-1}L_{FP}W(x,t')dt' \\ &= \Phi_t L_{FP}W(x,t), \end{aligned} \tag{27}$$

where

$$\Phi_t f(t) = \frac{\beta}{\Gamma(\alpha+1)}\frac{\partial}{\partial t}\int_0^t t'^{\beta/\alpha-1}(t^{\beta/\alpha}-t'^{\beta/\alpha})^{\alpha-1}f(t')dt'. \tag{28}$$

When $\beta = \alpha$, Eq.(27) reduces into the time fractional FPE (16). It is worthy to note that Mura et al. derived the integral form of Eq.(27) from a non-Markovian forward drift equation and called it stretched time-fractional diffusion equation [43].

The MSD $\langle x^2\rangle(t)$ of $Y(t) = x(t^\beta Z^{-\alpha})$ in a free force field can be computed by

$$\begin{aligned} \frac{d}{dt}\langle x^2\rangle(t) &= \int_{-\infty}^\infty x^2\frac{\partial W(x,t)}{\partial t}dx \\ &= \frac{2\beta}{\Gamma(\alpha+1)}\frac{d}{dt}\int_0^t t'^{\beta/\alpha-1}(t^{\beta/\alpha}-t'^{\beta/\alpha})^{\alpha-1}dt' \\ &= \frac{2\beta}{\Gamma(\alpha+1)}\frac{d}{dt}\int_0^{t^{\beta/\alpha}}\frac{\alpha}{\beta}(t^{\beta/\alpha}-s)^{\alpha-1}ds \\ &= \frac{2\beta}{\Gamma(\alpha+1)}t^{\beta-1}. \end{aligned} \tag{29}$$

So, we have

$$\langle x^2\rangle(t) = \frac{2}{\Gamma(\alpha+1)}t^{\beta}, \quad 0<\alpha<1, 0<\beta<2. \tag{30}$$

By comparing Eq.(30) with Eq.(18), we conclude that the diffusive process $Y(t)$ is controlled by parameter β in subordinator $s(t)$. It is easy to obtain that the subordinated process $Y(t)$ is subdiffusive when $0<\beta<1$, normal diffusive when $\beta=1$ and supdiffusive when $1<\beta<2$.

4 The subordinated process with subordinator $s(t) = t^{\beta}Z^{-\beta}$

Here we assume that $x(\tau)$ satisfies Langevin equation (6) and $s(t)$ has the form $s(t) = t^{\beta}Z^{-\beta}(0<\beta<2)$ where Z is an $\alpha-$stable random variable $(0<\alpha<1)$.

Since

$$P(s(t)\le\tau) = P(Z\ge\frac{t}{\tau^{1/\beta}}) = \int_{\frac{t}{\tau^{1/\beta}}}^{\infty}\psi_{\alpha}(t')dt', \tag{31}$$

we have

$$g(\tau,t) = \frac{1}{\beta}\frac{t}{\tau^{1/\beta+1}}\psi_{\alpha}(\frac{t}{\tau^{1/\beta}}). \tag{32}$$

After using the formulas (10) and (11), one can get the Laplace transform of the PDF $g(\tau,t)$

$$\widetilde{g}(\tau,u) = \frac{\alpha}{\beta}\tau^{\alpha/\beta-1}u^{\alpha-1}\exp\{-u^{\alpha}\tau^{\alpha/\beta}\}. \tag{33}$$

It is worth to note that the subordinated process $Y(t)$ constructed in this section can be connected to a CTRW with correlated waiting times (see Ref.[21]) in terms of Eq.(33). Indeed, in Ref.[21], the authors modeled CTRW with correlated waiting times by the following coupled Langevin equations

$$\frac{dx(\tau)}{d\tau} = \Gamma(\tau), \qquad \frac{dt(\tau)}{d\tau} = \int_0^{\tau}M(\tau-\tau')\eta(\tau')d\tau', \tag{34}$$

where $M(\tau)$ is a non-negative continuous memory function and $\eta(\tau)$ is a fully skew $\alpha-$stable Lévy noise. The Laplace transform of the PDF $g(\tau,t)$ of the process $s(t)$ defined by Eq.(5) has the form

$$\widetilde{g}(\tau,u) = u^{\alpha-1}\phi'(\tau)\exp\{-u^{\alpha}\phi(\tau)\}, \tag{35}$$

where $\phi(\tau) = \int_0^{\tau}d\tau'(\int_{\tau'}^{\tau}M(\tau''-\tau')d\tau'')^{\alpha}$. By comparing Eq.(35) with Eq.(33), we have $\phi(\tau) = \tau^{\alpha/\beta}$, which gives the connection between the subordinated process $Y(t)$ constructed in this section and the CTRW with correlated waiting times introduced in Ref. [21].

Some other kinds of correlated CTRW processes were given in Refs. [44-48]. The CTRW with correlated jumps was studied in Ref.[44]. The CTRWs with correlated waiting times defined as $t(\tau) = \int_0^{\tau}|L_{\alpha}(\tau')|d\tau'$ in Refs. [45-47] and with heterogeneous memorized waiting times defined as $D_{0+}^{1-\mu}t_x(\tau) = |x|^{\theta/\gamma}|L_{\alpha}(\tau)|$ in Ref. [48] are different from that introduced in Ref. [21].

Let us turn to the subordinated process $Y(t)$. In Laplace space, the PDF $W(x,t)$ can be represented by

$$\widetilde{W}(x,u) = \int_0^{\infty}f(x,\tau)\widetilde{g}(\tau,u)d\tau. \tag{36}$$

Inserting Eq.(33) into Eq.(36), we have

$$\begin{aligned}\widetilde{W}(x,u) &= \int_0^\infty f(x,\tau)\tfrac{\alpha}{\beta}\tau^{\alpha/\beta-1}u^{\alpha-1}\exp\{-u^\alpha\tau^{\alpha/\beta}\}d\tau \\ &= u^{\alpha-1}\int_0^\infty f(x,s^{\beta/\alpha})e^{-u^\alpha s}ds \\ &= u^{\alpha-1}\int_0^\infty f_1(x,s)e^{-u^\alpha s}ds \\ &= u^{\alpha-1}\widetilde{f}_1(x,u^\alpha),\end{aligned} \tag{37}$$

where $f_1(x,\tau)=f(x,\tau^{\beta/\alpha})$. Obviously there exists the relation

$$\frac{\partial}{\partial\tau}f_1(x,\tau)=\frac{\partial f(x,\tau_1)}{\partial\tau_1}\cdot\frac{d\tau_1}{d\tau}, \tag{38}$$

where $\tau_1=\tau^{\beta/\alpha}$.

Since the PDF $f(x,\tau)$ satisfies Eq.(7), we have

$$\begin{aligned}\tfrac{\partial}{\partial\tau}f_1(x,\tau) &= \tfrac{\partial f(x,\tau_1)}{\partial\tau_1}\cdot\tfrac{d\tau_1}{d\tau} \\ &= L_{FP}f(x,\tau_1)\cdot\tfrac{d\tau_1}{d\tau} \\ &= L_{FP}f_1(x,\tau)\tfrac{\beta}{\alpha}\tau^{\beta/\alpha-1}.\end{aligned} \tag{39}$$

We Set $q(\tau)=\frac{\beta}{\alpha}\tau^{\beta/\alpha-1}$, whose Laplace transform is $\widetilde{q}(u)=\frac{\Gamma(\beta/\alpha+1)}{u^{\beta/\alpha}}$. Taking Laplace transform for variable τ in Eq.(39), one gets

$$\begin{aligned}u\widetilde{f}_1(x,u)-f_1(x,0) &= \int_0^\infty L_{FP}f_1(x,\tau)q(\tau)e^{-u\tau}d\tau \\ &= \tfrac{1}{2\pi i}\int_{a-i\infty}^{a+i\infty}L_{FP}\widetilde{f}_1(x,s)\widetilde{q}(u-s)ds,\end{aligned} \tag{40}$$

with $0<a<Re(u)$.

From Eq.(37), with the initial condition $W(x,0)=f_1(x,0)=\delta(x)$, we have

$$\begin{aligned}u\widetilde{W}(x,u)-W(x,0) &= u^\alpha\widetilde{f}_1(x,u^\alpha)-f_1(x,0) \\ &= \tfrac{1}{2\pi i}\int_{a-i\infty}^{a+i\infty}L_{FP}\widetilde{f}_1(x,s)\widetilde{q}(u^\alpha-s)ds \\ &= \tfrac{1}{2\pi i}\int_{a-i\infty}^{a+i\infty}L_{FP}\widetilde{W}(x,s^{1/\alpha})s^{\frac{1-\alpha}{\alpha}}\tfrac{\Gamma(\beta/\alpha+1)}{(u^\alpha-s)^{\beta/\alpha}}ds \\ &= \tfrac{1}{2\pi i}\int_{a-i\infty}^{a+i\infty}s^{\frac{1-\alpha}{\alpha}}\tfrac{\Gamma(\beta/\alpha+1)}{(u^\alpha-s)^{\beta/\alpha}}ds\int_0^\infty L_{FP}W(x,t')e^{-s^{1/\alpha}t'}dt'.\end{aligned} \tag{41}$$

Setting $\widetilde{k}(u,s)=s^{\frac{1-\alpha}{\alpha}}\frac{\Gamma(\beta/\alpha+1)}{(u^\alpha-s)^{\beta/\alpha}}$ and taking inverse Laplace transform for $u\to t$ in Eq.(41), we obtain

$$\frac{\partial}{\partial t}W(x,t)=\Phi_t L_{FP}W(x,t), \tag{42}$$

where

$$\Phi_t h(t) = \frac{1}{2\pi i}\int_{a-i\infty}^{a+i\infty} k(t,s)ds \int_0^\infty h(t')e^{-s^{1/\alpha}t'}dt'. \tag{43}$$

Next we also compute the MSD $\langle x^2\rangle(t)$ of the subordinated process $Y(t) = x(t^\beta Z^{-\beta})$ in a free force filed.

Since

$$\langle x^2\rangle(t) = 2\int_0^\infty \tau g(\tau,t)d\tau, \tag{44}$$

we have

$$\begin{aligned}\langle x^2\rangle(u) &= 2\textstyle\int_0^\infty \tau\widetilde{g}(\tau,u)d\tau \\ &= 2\textstyle\int_0^\infty \tau\frac{\alpha}{\beta}\tau^{\alpha/\beta-1}u^{\alpha-1}\exp\{-u^\alpha\tau^{\alpha/\beta}\}d\tau \\ &= 2u^{\alpha-1}\textstyle\int_0^\infty s^{\beta/\alpha}e^{-u^\alpha s}ds \\ &= \frac{2\Gamma(\frac{\beta+\alpha}{\alpha})}{u^{\beta+1}}.\end{aligned} \tag{45}$$

So

$$\langle x^2\rangle(t) = \frac{2\Gamma(\beta/\alpha)}{\alpha\Gamma(\beta)}t^\beta. \tag{46}$$

From Eq.(46), we conclude that the subordinated process is subdiffusive when $0<\beta<1$, normal diffusive when $\beta=1$ and supdiffusive when $1<\beta<2$.

5 Conclusions

In this work, we consider the subordinated process $Y(t)$ which is controlled by a family of subordinators. For the case of $s(t) = t^\alpha Z^{-\alpha}$, the PDF $W(x,t)$ of the subordinated process $Y(t)$ satisfies a time-fractional Fokker-Planck equation. For the case of $s(t) = t^\beta Z^{-\alpha}$, the corresponding diffusion equation is a stretched time-fractional Fokker-Planck equation. For the case of $s(t) = t^\beta Z^{-\beta}$, we obtain that the Laplace transform of the PDF $g(\tau,t)$ is of the form $\widetilde{g}(\tau,u) = \frac{\alpha}{\beta}\tau^{\alpha/\beta-1}u^{\alpha-1}\exp\{-u^\alpha\tau^{\alpha/\beta}\}$. The resulting process can be connected with CTRW with correlated waiting time.

Acknowledgments

This project was supported by the Natural Science Foundation of China (Grant Nos. 11371016 and 11271311), the Chinese Program for Changjiang Scholars and Innovative Research Team in University (PCSIRT) (Grant No. IRT1179), the Lotus Scholars Program of Hunan province of China.

References

[1] J.P.Bouchaud, A.Georges, Anomalous diffusion in disordered media: statistical mechanisms models and physical applications, Phys. Rep. 195 (1990) 127-293.

[2] J.Klafter, I.M.Sokolov, Anomalous diffusion spreads its wings, Physics World, August (2005) 29-32.

[3] I.Eliazar, J.Klafter, Anomalous is ubiquitous, Anna. Phys. 326 (2011) 2517-2531.

[4] R.Metzler, J.Klafter, The random walks guide to anomalous diffusion: a fractional dynamics approach, Phys. Rep. 339 (2000) 1-77.

[5] E.W.Montroll, G.H.Weiss, Random walks on lattices.II, J. Math. Phys. 6 (1965) 167-181.

[6] H.C.Fogedby, Langevin equations for continuous time Lévy flights, Phys. Rev. E 50 (1994) 1657-1660.

[7] S.Bochner, Subordination of non-Gaussian stochastic processes, Proc. Natl. Acad. Sci. USA 48 (1962) 19-22.

[8] Metzler R and Klafter J, 2000 *J. Phys. Chem.* B **104** 3851

[9] Metzler R and Klafter J, 2000 *Phys. Rev.* E **61** 6308

[10] Metzler R, 2000 *Phys. Rev.* E **62** 6233

[11] A.Piryatinska, A.I.Saichev, W.A.Woyczynski, Models of anomalous diffusion: the subdiffusive case, Phys. A 349 (2005)375-420.

[12] A.Baule, R.Friedrich, Joint probability distributions for a class of non-Markovian processes, Phys. Rev. E 71 (2005) 026101.

[13] M.Magdziarz, A.Weron, K.Weron, Fractional Fokker-Planck dynamics: Stochastic representation and computer simulation, Phys. Rev. E 75 (2007) 016708.

[14] M.Magdziarz, A.Weron, Competition between subdiffusion and Lévy flights: A Monte Carlo approch, Phys. Rev. E 75 (2007) 056702.

[15] T.Koren, J.Klafter, M.Magdziarz, First passage times of Lévy flights coexisting with subdiffusion, Phys. Rev. E 76 (2007) 031129.

[16] D.Kleinhans, R.Friedrich, Continuous-time random walks: Simulation of continuous trajectories, Phys. Rev. E 76 (2007) 061102.

[17] R.Gorenflo, F.Mainardi, A.Vivoli, Continuous-time random walk and parametric subordination in fractional diffusion, Chaos, Solitons and Fractals 34 (2007) 87-103.

[18] A.Weron, M.Magdziarz, K.Weron, Modeling of subdiffusion in space-time-dependent force fields beyond the fractional Fokker-Planck equation, Phys. Rev. E 77 (2008) 036704.

[19] M.Magdziarz, A.Weron, J.Klafter, Equivalence of the fractional Fokker-Planck and subordinated Langevin equations: the case of a time-dependent force, Phys. Rev. Lett. 101 (2008) 210601.

[20] M.Magdziarz, Stochastic representation of subdiffusion processes with time-dependent drift, Stoch. Proc. Appl. 119 (2009) 3238-3252.

[21] A.V.Chechkin, M.Hofman, I.M.Sokolov, Continuous-time random walk with correlated waiting times, Phys. Rev. E 80 (2009) 031112.

[22] S.Eule, R.Friedrich, Subordinated Langevin equations for anomalous diffusion in external potentials - Biasing and decoupled external forces, EPL 86 (2009) 30008.

[23] B.Dybiec, E.Gudowska-Nowak, Subordinated diffusion and continuous time random walk asymptotics, Chaos 20 (2010) 043129.

[24] R.Gorenflo, F.Mainardi, Subordination pathways to fractional diffusion, Eur. Phys. J. Special Topics 193 (2011) 119-132.

[25] L.Lv, W.Qiu, F.Ren, Fractional Fokker-Planck equation with space and time dependent drift and diffusion, J. Stat. Phys. 149 (2012) 619-628.

[26] M.Magdziarz, Langevin picture of subdiffusion with infinitely divisible waiting times, J. Stat. Phys. 135 (2009) 763-772.

[27] J.Gajda, M.Magdziarz, Fractional Fokker-Planck equation with tempered $\alpha-$stable waiting times: Langevin picture and computer simulation, Phys. Rev. E 82 (2010) 011117.

[28] J.Janczura, A.Wyłomańska, Anomalous diffusion models: different types of subordinator distribution, Acta Phys. Pol. B 43 (2012) 1001-1016.

[29] A.Wyłomańska, Arithmetic Brownian motion subordinated by tempered stable and inverse tempered stable processes, Phys. A 391 (2012) 5685-5696.

[30] J.Gajda, Fractional Fokker-Planck equation with space dependent drift and diffusion: the case of tempered $\alpha-$stable waiting times, Acta Phys. Pol. B 44 (2013) 1149-1161.

[31] Y.X.Zhang, H.Gu, J.R.Liang, Fokker-Planck type equations accociated with subordinated processes controlled by tempered $\alpha-$stable processes, J. Stat. Phys. 152 (2013) 742-752.

[32] Teuerle M, Wyłomańska A and Sikora G, 2013 *J. Stat. Mech.* P05016

[33] A.Wyłomańska, The tempered stable process with infinitely divisible inverse subordinators, J. Stat. Mech. (2013) P10011.

[34] J.Gajda, A.Wyłomańska, Fokker-Planck type equations associated with fractional Brownian motion controlled by infinitely divisible processes, Phys. A 405 (2014) 104-113.

[35] M.M.Meerschaert, H.P.Scheffler, Limit theorems for continuous-time random walks with infinite mean waiting time, J. Appl. Prob. 41 (2004) 623-638.

[36] H.Risken, The Fokker-Planck Equation, Springer-Verlag Berlin Heidelberg, New York, 1989.

[37] R.Metzler, E.Barkai, J.Klafter, Anomalous diffusion and relaxation close to thermal equilibrium: a fractional Fokker-Planck equation approach, Phys. Rev. Lett. 82 (1999) 3563-3567.

[38] R.Metzler, E.Barkai, J.Klafter, Anomalous transport in disordered systems under the influence of external fields, Phys. A 266 (1999) 343-350.

[39] E.Barkai, Fractional Fokker-Planck equation, solution, and application, Phys. Rev. E 63 (2001) 046118.

[40] I.Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.

[41] R.Metzler, E.Barkai, J.Klafter, Driving fractional Fokker-Planck equations from a generalised master equation, EPL 46 (1999) 431-436.

[42] E.Barkai, R.Metzler, J.Klafter, From continuous time random walks to the fractional Fokker-Planck equation, Phys. Rev. E 61 (2000) 132-138.

[43] A.Mura, M.S.Taqqu, F.Mainardi, Non-Markovian diffusion equations and processes: Analysis and simulations, Phys. A 387 (2008) 5033-5064.

[44] Meerschaert M M, Nane E and Xiao Y, 2009 *Stat. Probab. Lett.* **79** 1194

[45] Tejedor V and Metzler R, 2010 *J. Phys.* A **43** 082002

[46] Magdziarz M, Metzler R, Szczotka W and Zebrowski P, 2012 *J. Stat. Mech.* P04010

[47] Magdziarz M, Metzler R, Szczotka W and Zebrowski P, 2012 *Phys. Rev.* E **85** 051103

[48] Wang J, Zhou J, Lv L J, Qiu W Y and Ren F Y, 2014 *J. Stat. Phys.* **156** 1111